NASA Technical Memorandum 103263

Comment on "Local Energy Transfer and Nonlocal Interactions in Homogeneous, Isotropic Turbulence" [Phys. Fluids A2, 413 (1990)]

Robert G. Deissler

Lewis Research Center

Cleveland, Ohio

August 1990

(NASA-TM-103263) COMMENT ON LOCAL ENERGY TRANSFER AND NONLOCAL INTERACTIONS IN HOMOGENEOUS, ISOTROPIC TURBULENCE (PHYS. FLUIDS AZ, 413 (1990)) (NASA) 7 p CSCL 20D

N91-13638

Unclas 0308330





COMMENT ON "LOCAL ENERGY TRANSFER AND NONLOCAL INTERACTIONS IN HOMOGENEOUS, ISOTROPIC TURBULENCE"

[PHYS. FLUIDS A2, 413 (1990)]

Robert G. Deissler National Aeronautics and Space Administration Lewis Research Center Cleveland, Ohio 44135

ABSTRACT

It is argued that the low-Reynolds number results in the subject paper are not congruous with the data of Ling and Huang (Phys. Fluids $\underline{13}$, 2912 (1970)), and that the spectral transfer should be less local.

Domaradzki and Rogallo¹ address an important question concerning the nature of the spectral transfer in turbulence: Does the transfer take place mainly between neighboring wavenumbers, or between wavenumbers that are substantially separated? In spite of the fundamental nature of this question, only a limited amount of work has appeared in the literature.

My comments have to do with the correctness of certain results in the subject paper, and with that of my results in Ref. 2. The results under question concern contributions $P(\kappa,\kappa')$ from wavenumbers κ' to the total energy transfer $T(\kappa)$ at κ , where $T(\kappa)$ was measured by Ling and Huang.³ The authors of the subject paper believe that the results in Ref. 2 are in error because of the disagreement of those results with their own, and because the method used there does not give unique results for $P(\kappa,\kappa')$.

In reality, however, the conditions used by the writer to obtain $P(\kappa,\kappa')$ from the $T(\kappa)$ measured by Ling and Huang³ are quite restrictive. Those conditions are⁴

$$P(\kappa,\kappa') = -P(\kappa',\kappa) \tag{1}$$

and

$$T(\kappa) = \int_0^\infty P(\kappa, \kappa') d\kappa'. \qquad (2)$$

As will be seen, the present method,² in contrast to the numerical simulation of the subject paper, gives results for $P(\kappa,\kappa')$ which satisfy Eqs. (1) and (2) exactly when the empirical equation of Ling and Huang is used for $T(\kappa)$. That equation, which gives an excellent fit to Ling and Huang's data is, in dimensionless form,

$$T(\kappa) = -\frac{10}{3} \left(\kappa + \sqrt{10}\kappa^2 + 3\kappa^3 - 2\sqrt{10}\kappa^4\right) e^{-\sqrt{10}\kappa}$$
(3)

where κ has been scaled by setting $\left[v(t-t_0)\right]^{1/2}\kappa=\kappa$, T is a dimensionless

transfer function, ν is the kinematic viscosity, t is the time, and t₀ is the virtual origin of the turbulence. Equation (3) is plotted in Fig. 1.

The description of the method for obtaining $P(\kappa,\kappa')$ which follows is perhaps clearer and more satisfying than that given in Ref. 2. It will be seen that, for the low Reynolds-number data, $P(\kappa,\kappa')$ can be obtained by using a reasonable systematic procedure.

Terms in the expression for $P(\kappa,\kappa')$ that satisfy Eq. (1), and, through Eq. (2), can satisfy Eq. (3), are of the form

Term =
$$(\kappa^m \kappa^{,n} - \kappa^n \kappa^{,m}) e^{-\sqrt{10}(\kappa + \kappa^{,n})},$$
 (4)

where m and n are positive intergers. Then the expression for $P(\kappa,\kappa')$ which satisfies Eqs. (1) to (4), while containing the smallest possible number of terms of the lowest possible order in κ and κ' , is

$$P(\kappa,\kappa') = \frac{100}{3} \left[\sqrt{10} (\kappa \kappa'^2 - \kappa^2 \kappa') + 3(\kappa \kappa'^3 - \kappa^3 \kappa') - 2\sqrt{10} (\kappa \kappa'^4 - \kappa^4 \kappa') \right] e^{-\sqrt{10}(\kappa + \kappa')}$$
(5)

Note that a term proportional to $\kappa - \kappa'$ is not included in Eq. (5), since that would not satisfy T(0) = 0, as required by Eq. (3), although it would satisfy Eq. (1).

An important consequence of Eq. (2) and the relation $P(\kappa,\kappa') = -P(\kappa',\kappa)$ (Eq. (1)) is that if $T \propto \kappa^n$ for small κ , then $P \propto \kappa'^n$ or $P \propto \kappa^n$ for κ' or κ respectively small. In the present case this means that $T(\kappa)$ and $P(\kappa,\kappa')$ both start with the first power of κ or κ' (see Eqs. (3) and (5), or Figs. 1 and 2). This has a major effect on the shape of $P(\kappa,\kappa')$ and tends to make the bulk of the energy transfer from κ' to κ nonlocal. (See the solid curve in Fig. 2, where the peak occurs at small values of κ'/κ .) On the other

hand, the shape of the curve obtained in the subject paper near $\kappa'=0$ (dashed curve in Fig. 2) is characteristic of that of a curve which starts with a much higher power of κ' than the first. It thus appears to be not congruous with the energy transfer of Ling and Huang (Eq. (3)) and conditions (1) and (2). As a result, the dashed curve shows an energy transfer which is evidently too local (the peak of the curve lies too close to $\kappa'/\kappa=1$). The simulation from the subject paper does not seem to faithfully represent the experiment of Ling and Huang. The present representation of $P(\kappa,\kappa')$, in starting with the first power of κ' , appears to represent the experiment much more closely. Thus, it appears that the original conclusion that the spectral transfer in Ling and Huang's experiment is nonlocal (the energy tends to jump between wavenumbers that are separated) still holds.

As further evidence of the importance of the power of κ' with which $P(\kappa,\kappa')$ starts, in determining the localness of the spectral transfer, we note that in Ref. 5, where a different method is used to determine $P(\kappa,\kappa')$, and where P starts with a power of κ' much higher than the first, the transfer is more local (closer to that in the subject paper). However, when $P(\kappa,\kappa')$ was assumed to start with the first power of κ' , as in the experiment of Ling and Huang, the transfer was, as expected, less local, and closer to that given by Eq. (5).

As a final note it might be pointed out that the spectral transfer must always be somewhat nonlocal because of the condition that P=0 for $\kappa'/\kappa=1$. That condition follows from Eq. (1) by letting $\kappa'=\kappa$. Thus the descrepency between the results of Domaradzki and Rogallo¹ and the present results² is one of degree, rather than a qualitative difference.

REFERENCES

- 1 J.A. Domaradzki and R.S. Rogallo, Phys. Fluids $\underline{A2}$, 413 (1990).
- 2R.G. Deissler, Appl. Sci. Res. 34, 379 (1978).
- 3 S.C. Ling and T.T. Huang, Phys. Fluids, $\underline{13}$, 2912 (1970).
- ⁴G.K. Batchelor, <u>The Theory of Homogeneous Turbulence</u> (Cambridge University Press, New York, 1953.
- 5R.G. Deissler, Phys. Fluids, $\underline{1}$, 111 (1958).




Figure 1.—Experimental energy transfer spectrum from Ling and Huang³ (Eq. (3)) for low and moderate Reynolds numbers (R_{λ} is between 3 and 30). κ_0 is the wave number where T(κ) = 0 ([ν (t - t₀)]^{1/2} κ_0 = 1.0758).




Figure 2.—Calculated contributions $P(\kappa, \kappa')$ to net energy transfer T at κ/κ_0 in figure 1 = 2.2.

National Aeronautics and Space Administration	Report Docum	nentation Pa	age	
1. Report No. NASA TM-103263	2. Government Acce	ession No.	3. Recipient's Catal	og No.
4. Title and Subtitle			5. Report Date	
Comment on "Local Energy Transfer and Nonlocal Interactions			August 1990	
Homogeneous, Isotropic Turk	bulence'' [Phys. Fluids A2,	413 (1990)]	6. Performing Organ	nization Code
7. Author(s)			Performing Organization Report No.	
Robert G. Deissler			E-5696	
			10. Work Unit No.	
A. Borforming Organization Name and	I Addrose		505-90-01	
9. Performing Organization Name and Address			11. Contract or Grant No.	
National Aeronautics and Spa Lewis Research Center	ice Administration			
Cleveland, Ohio 44135-319	1		13. Type of Report a	nd Period Covered
12. Sponsoring Agency Name and Address			Technical Memorandum	
National Aeronautics and Spa		14. Sponsoring Agend	cy Code	
Washington, D.C. 20546-00	JO I			
16. Abstract It is argued that the low-Rey	nolds number results in the s	ubiect paper are	not congruous with the	e data of Ling
and Huang (Phys. Fluids 13,	and that the s	poetrai transfer s	ACCURATE DE 1000 100011.	
17. Key Words (Suggested by Author(s))	18. Distribution St	atement	
Energy transfer Localness Turbulence Homogeneous		Unclassified – Unlimited Subject Category 34		
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of pages	22. Price*
19. Security Classif. (of this report) Unclassified	· · · · · · · · · · · · · · · · · · ·	lassified	6	A02